



# ORTHOPEDIC ASSOCIATES of HARTFORD

## Achilles Tendon Rupture Repair Return to Sport Protocol

### Keys for Achilles Tendon Rehabilitation:

- Calf girth, Plantar Flexor strength and ankle ROM are all keys to full recovery. The ideal is to achieve full symmetry in all 3 areas.
  - ROM should be measure in both Open and Closed-Chain methods
  - Strength can be measured through use of the Single Heel Rise test and MMT should be scored using the criteria below:

### Testing Plantar Flexion in Standing:

- Grading:
    - **5/5 (N)**= Patient successfully raises heel from floor through entire ROM of plantar flexion. Patient should complete a minimum of 20 repetitions in good form and without apparent fatigue. (tibialis posterior and the peroneus (fibularis) longus and brevis muscles must be between a 4/5 and 5/5 to stabilize the forefoot and provide counter pressure against the floor).
      - \*20 heel rises represents over 60% of maximum electromyographic activity of the plantar flexors
    - **4/5 (G)**= Patient can complete 10-19 repetitions of heel rises with full ROM and then has difficulty completing the movement.
    - **3/5 (F)**= Full ROM for 1-9 repetitions of heel rises, and becomes fatigued.
    - **2+/5 (P)**= Patient can just clear the heel from the floor and cannot get up on the toes for full ROM.
    - **No grade for 2/5 in the standing position, all others tested in prone position with feet off the edge of the table.**
    - **2+/5 (P+)**= patient is able to plantar flex through available ROM against maximal resistance.
    - **2/5 (P)**= Patient is able to complete plantar flexion full ROM however cannot tolerate resistance.
    - **2-/5 (P-)**= Patient is able to complete only partial ROM.
    - **1/5 (T)**= Their tendon reflects some contractile activity in the muscle, however, no joint motion occurs. Contractile activity may be palpated in the muscle bellies. (The Gastrocnemius at mid-calf with thumb and finger on either side of the midline, above the soleus. Soleus is best palpated distally on the posterolateral surface).
    - **0/5 (ZERO)**= no contractile activity
      - \* If the patient is unable to lie prone, alternate test position is supine, clinician cupping the heel, with their forearm against the plantar surface of the patient's foot, and a force applied into Dorsiflexion, **after full ROM was assessed.**
- \*Taken from Daniels and Worthington (2002) Muscle Testing: Techniques of Manual Examination, 5<sup>th</sup> edition.**

## Postoperative Achilles Tendon Rupture Repair Return to Sport Protocol

### Immediately Post-Op- Protection Phase (0-4 weeks Post-Op)

#### Goals:

- Protect healing tissue
- Decrease/eliminate inflammation and edema, while controlling pain
- Prevent muscle atrophy and wound infection (to minimize scar formation)

#### Patient Education:

- Posterior Splint or Boot must be worn day and night.
- Incision must be kept clean and dry.
- Non-Weight-Bearing for 3 weeks post-op, ambulation with bilateral crutches
- Cold Packs /Cryocuff should be used multiple times through the day with the leg elevated for first 1-2 weeks, then PRN

#### Treatments:

- Splint/Brace:
  - Posterior splint in 10°-20° plantar flexion 24 hours/day for first 3 weeks, Progressing to boot at ~week 3 post-op (to be wore awake and while sleeping) with heel wedge to keep at ~10° plantar flexion.
- Weight-Bearing:
  - Non-Weight-Bearing for first 3 weeks Post-op, then patient may begin partial Weight-Bearing with Bilateral crutch in the boot with heel lift, (depending on MD Assessment).
- Modalities:
  - Cold Packs post treatment to control pain, swelling, and inflammation
  - Moist Hot Packs are not to be utilized until 2 weeks post-op
- Gait Training:
  - Gait and stair training with crutches, for proper technique
- Manual Therapies & ROM:
  - PROM: Clinician administered PROM is not to be performed for 8 Weeks post-op, Patient should perform Gravity-Assisted PROM into plantar flexion in a seated position (once incision is healed).
  - AROM: ~3 weeks post-op patient may begin AROM exercises of the ankle, **DF to neutral only with bent knee position, no active PF**
- Proximal LE and Upper Body Kinetic Chain Strengthening Exercises:
  - **No Ankle Strengthening** for 1st 2 weeks, Sub-max. after 2 weeks no PF
  - 4-Way Hip SLR (Flex, Ext, Abd, Add), H/S curls
  - Foot Intrinsic Strengthening: Towel crutches, marbles, toe extension, etc.
  - Any UE and Core exercises in a LE non-weight-bearing position
  - Stationary Bicycle in the Boot

### Intermediate Protection Phase: (4-8 Weeks)

#### Goals:

- Continue to protect the healing Achilles repair
- Continue to decrease/eliminate pain and inflammation
- Continue to control/prevent muscular atrophy
- Progress to full weight-bearing
- Regain ankle range of motion (0°/neutral DF)

### **Patient Education:**

- Educate patient on necessity to continue to wear the walking boot and adhere to restrictions to continue to protect the Achilles repair while early motion and weight-bearing is initiate/progressed.
- Brace must be worn whenever patient is in weight-bearing, it may be removed for shower/bath and to perform exercises.

### **Treatments:**

- Weight-Bearing:
  - Progress WBAT with Boot and 1-inch heel lift for 2 weeks, and then decrease to ½ inch heel lift, D/C crutches when full weight-bearing in boot is achieved.
- Modalities:
  - Continue all modalities PRN for pain, inflammation, and swelling
- Gait Training:
  - Gait Training progression from PWB/WBAT@WB as tolerated.
- Manual Therapies:
  - Gentle scar mobilization to Achilles
  - PROM may be initiated at 8 weeks post-op, gentle into DF initially
- Exercises:
  - Continue foot strengthening, SLR exercises, leg press (no PF movement)
  - Seated Baps Board, Ankle gentle AROM- (alphabets, rotations, pumping)
  - End-range Isometrics, supine/ seated calf raises (low/no resist)
  - **Weight Shifting:** Towel roll under heels @ 4 weeks (no boot)
    - Bilateral stance with boot, progress to stagger stance –week 5
      - Continue Stationary Bicycle with boot
      - Continue Upper Body& Core exer. (seated/lying/CKC w/boot/LE on ball)

### **Criteria for progression to Initial Strengthening phase:**

- Full Weight-Bearing with Boot out of home, No Boot in home and clinic
- No Increased symptoms with current AROM & PROM exercises
- 0°-5° Dorsiflexion/ Neutral ankle ROM achieved- to be able to D/C boot
- Swelling/Edema controlled

### **Initial Strengthening Phase (~8-12 Weeks)**

#### **Goals:**

- Increase ROM to 10° Dorsiflexion, Achieve full ankle ROM INV, EV, PF
- Continue to progress weight bearing tolerance, D/C Walking Boot all surfaces (~10 weeks)
- Normalized Gait
- Restore normal joint arthrokinematics of all foot and ankle joints
- Continue to protect healing tissues
- Decrease calf atrophy

### **Patient Education:**

- Education on progressing weight-bearing slowly, based on ROM and tolerance.
- Patient should continue to wear the boot, brace and/or heel lift until MD D/C's

## Treatments:

- Weight Bearing:
  - Continue to progress weight bearing to FWB'ing out of boot (with 1/4-1/2 inch heel lift in shoe/sneaker)~ 10 weeks post-op
- Gait Training: (MD may prescribe an ankle brace once boot is D/C'd).
  - ~10 Weeks taper out of boot in the home and clinic, continue to wear out of the home. ~12 weeks D/C boot if appropriate (**MD will D/C boot**).
- Manual Therapies:
  - Manual Mobilization of foot & Midfoot, gentle Subtalar & Talo-Crural mobilization- to help restore foot and ankle ROM
  - PROM all planes to improve foot and ankle mobility
- Exercises:
  - Light resistance band 4- way ankle exercise- gradually increase resistance
  - Seated (light resist), Standing (PWB) BAPS & decline boards- all planes
  - Forward, Retro and Lateral step up & downs- low step height
  - Progress DL heel raises supine, seated & add standing as tol. (~12 weeks)
  - Continue LE, Foot intrinsic, Hip, Core, & UE exercises, & Leg Press
  - Continue LE bicycle, can begin Retro Treadmill walking

## End of Phase Testing:

- Core Testing: (**See attached sheets for description**): 1-Segmental Multifidus Test, 2-Trunk Curl Up Test, 3-Double-Leg Lowering Test, 4-Side Bridge Test, 5-Prone Bridge Test, 6-Supine Single-Leg Bridge Test, 7-Extensor Endurance Test

## Criteria for progression to Advanced Strengthening phase:

- $\geq 10^\circ$  Ankle Dorsiflexion, all other planes in normal ROM
- No pain, inflammation or swelling
- Normal Gait
- No increased symptoms with any of the current exercises

## Advanced Strengthening Phase (~12-22 Weeks)

### Goals:

- Progress to full ankle ROM in all planes
- Increase strength and balance to be able to begin advanced activity skills
- No calf/muscle atrophy
- D/C brace if still being used (**by MD only**)
- Get Achilles Tendon Total Rupture Score (ATRS)- Self -Reported Patient Evaluation
- Good proprioception/balance in single-leg stance
- Gradually introduce sub-maximal agility & reaction drills (~18-20 weeks post-op), and sport-like activities (20-22 weeks post-op)
- Get baseline FMS™ Assessment Screen & Y-Balance Test™ (~12-14 weeks)
- Initiate Walk-to Run (~16-18 weeks post-op), LE Plyometric Protocols (~18-20 weeks post-op)

## Treatments:

- Continue to progress Dorsiflexion to normal ROM
- Continue to increase repetitions and resistance: As tolerated
  - 4-way ankle resistance band exercises
  - Standing DL calf raises, add eccentric lowering; SL toward end of phase
    - Begin Heel Raise progression protocol (**See Attached Sheet**)
  - Anterior, Lateral, Retro step-ups & downs- increasing step height
- Continue to advance Core, Hip, and UE stabilization exercises
- Teach Injury Prevention/ Athletic Dynamics Stretching exercises- (See Sheets)
- Squats: DL (week 12), SL (~week 16)
  - 0-35° 12 weeks post-op
  - 0-60° 14 weeks post-op
- Retro treadmill walking
- Continue leg press and add calf press raises- DL and SL
- Anterior posterior and lateral lunges
- Begin Walk-to Run Protocol (~16-18 weeks post-op)- when SL Squat to 60° knee flexion with no knee valgus or pain in LE, especially Achilles
- Begin LE Plyometric Protocol (~week18 post-op)
- Begin light sub-maximal Agility and Reaction drills, Sport-like activities (~18-20 weeks post-op)
  - Straight line drills, ladder drills, step over cone drills, skipping, hopping

## End of Phase Testing:

- FMS™ and Y-Balance Test™ LE score
  - FMS® (goal of  $\geq 14/21$  points with no 0/3, 1/3, or asymmetries) and Y-Balance™ score (Goal- Statistically symmetrical to non-injured leg)
- Perform hop testing: See Attached Sheets
  - 2 practice trials, 2 timed/measured trials; average injured to non-injured
  - 1- Single-leg hop for distance
  - 2- Triple hop for distance
  - 3- Single-Leg Crossover triple hop
  - 4- 6-Metered timed hop
  - Hop-To-Stop Jump Tests: SL and SL with Opposite Landing- 3 trials of each bilateral
  - Tuck Jump- 2 trials for 10 seconds each (**See Scoring Criteria**)
  - DL Jump Test- Modified with hands behind back- (**See Scoring Criteria**)
- SL Squat Test: (0°-60°):
  - No knee valgus or ankle instability noted
- Dorsiflexion Lunge Test (bilateral)
- Core Testing: Re-Test (Goal- No major dysfunction/ strength deficits noted)
  - Segmental Multifidus Test
  - Trunk Curl Up Test
  - Double-Leg Lowering Test
  - Side Bridge Test
  - Prone Bridge Test

- Supine Single-Leg Bridge Test
- Extensor Endurance Test

### **Criteria to progress to Return to Activity Phase**

- 1- Full ROM
- 2- No pain or inflammation
- 3- A score of  $\geq 14/21$  on the FMS™ Assessment Screen, with **No 0/3= pain** on any of the 7 fundamental movement patterns
- 4- No statistical asymmetries and on the Y-Balance Test™
- 5- Hop Tests (90% or higher compared to non-injured leg)
- 6- Limb Symmetry Index (LSI) of 90% or greater on hop tests
- 7- No major deficits noted with Core and SL Squat testing
- 8-  $\geq 25^\circ$  Dorsiflexion Lunge Test

### **Return to Activity Phase (~22-32 weeks)**

\* Patient may not still be in Rehab at this point on a regular basis, so home exercises program needs to be demonstrated, and explained in detailed (pictures and written).

- If Patient is continuing PT at home, it is expected that they are to be re-evaluated every 4-6 weeks for advancement of exercises as well as determining progression.

### **Goals:**

- Continue to increase to full LE strength
- Continue to increase power, endurance, agility, coordination, and balance
- Gradual return to sport activities
- Get Achilles Tendon Total Rupture Score (ATRS)- Self -Reported Patient Evaluation

### **Treatment:**

- Continue advanced strengthening, core, proprioceptive, and mobility exercises
- Continue/Finish Heel Raise Progression Protocol (**See Attached Sheet**)
- Continue Walk-to-Run Protocol- if not completed to running 2 miles
- Teach Sport Specific Dynamic Stretching routine/Warm-up (**See Sheets**)
- Complete LE Plyometric Protocol
  - Progress to Box Jumps
- Progress agility and reaction exercises- increase speed, distance, and perturbations
- Begin Interval Running Protocol- when cleared by MD to perform cutting and sprinting activities
- Begin Interval Sport Protocol (ex. Kicking, Throwing, etc.) based on the sport patient will be returning- (See Sheets)

**Functional Testing: for progression to sport-specific training/Return to Sport-\***Can be perform over multiple days- See Attached Sheets for descriptions of tests and norms for data

### **Strength and Power Testing:**

- Sargent Vertical Jump Test
- Figure-8 Hop Test
- Up-Down Test
- Hexagon Test (DL), Modified Hexagon Hop Test (SL)

### Speed, Agility, and Quickness Testing:

- Agility T-Test, or Modified Agility T-Test (MAT)
- Three-Cone Drill Test
- Slalom Test
- Backward Movement Agility Test
- Zigzag Run Test
- Lower Extremity Functional Test (LEFT)

### Core Testing:

- Segmental Multifidus Test
- Trunk Curl Up Test
- Double-Leg Lowering Test
- Side Bridge Test
- Prone Bridge Test
- Supine Single-Leg Bridge Test
- Extensor Endurance Test

### Hop/Jump Testing:

- 2 practice trials, 2 timed/measured trials; average injured to non-injured
- 1- Single-leg hop for distance
- 2- Triple hop for distance
- 3- Single-Leg Crossover triple hop
- 4- 6-Metered timed hop
- Hop-To-Stop Jump Tests: SL and SL with Opposite Landing- 3 trials of each bilateral
- Tuck Jump- 2 trials for 10 seconds each (**See Scoring Criteria Sheets**)
- DL Jump Test- Modified with hands behind back- 2 trials (**See Scoring Criteria Sheets**)

### Function, Balance, and Endurance Testing:

- FMS® Assessment Screen
- Y-Balance Test™
- Vail Sport Test

### **\*Return-to-Sport Activities**

#### Criteria for Return-to-Specific Sport Activities:

- **\*Ultimate decision is made by MD**
- Achilles Tendon Total Rupture Score (ATRS):  $\geq 95/100$
- A score of  $\geq 16/21$  on the FMS® Assessment Screen, with no pain or asymmetries, especially with Inline-lunge, and 2 or greater on all 7 fundamental movement patterns
- Y-Balance Test™- No asymmetries and a composite score as close to 100 as possible
- Limb Symmetry Index (LSI) of  $\geq 95\%$  on SL hop tests compared to Uninjured Leg
- LEFT Score:  $\leq 117$  seconds (females),  $\geq 105$  seconds (males)
- Vail Sport Test™:  $\geq 46/54$  points
- Isokinetic testing: **(if available)**
  - Quadriceps (90% or greater) compared to non-injured leg
  - Hamstring (100%-110%) compared to non-injured leg
  - Hamstring-Quadriceps Ratio (80% or greater)
- DL Jump and Tuck Jump: meets scoring criteria

- Sargent Vertical Jump: **See chart** for norms based on age of athlete
- Agility T-Test: Adults:  $\leq 8.9$ -13.5 seconds, Adolescents:  $\leq$
- Dorsiflexion Lunge Test: distance of foot to wall  $\geq 9$  cm, & tibial shaft angle  $\geq 35^\circ$
- No discomfort or swelling, and passing/statistically equal to normative values (if available) with above Functional Tests

### Heel Raise Progression Protocol

- **Phase 1:** Performed by patient **Daily- 3X/ Day**
  - 20 Repetitions per session
  - Technique: Double-Leg Heel Raises- Both heel Up and Down simultaneously
  - Duration 2-4 weeks
- **Phase 2:** Performed **5 days/ Week (2 days on, 1 Day off)- 1-2X/ Day**
  - Technique:
    - DL Heel Raise Up and Down together X 10 Repetitions
    - DL Heel Raise Up together, Raise Unaffected Leg and Down/Lower Affected Leg Only X 10 Repetitions
    - DL Heel Raise Up and Down Together X 10 Repetitions
  - Duration 2 weeks
- **Phase 3:** Performed **5 days/ Week (2 days on, 1 Day off)- 1-2X/ Day**
  - Technique:
    - DL Heel Raise Up and Down together X 10 Repetitions
    - DL Heel Raise Up together, Raise Unaffected Leg and Down/Lower Affected Leg Only X 10 Repetitions
    - DL Heel Raise Up and Down Together X 10 Repetitions
  - Duration 2 weeks
- **Phase 4:** Performed **5 days/ Week (2 days on, 1 Day off)- 1X/ Day**
  - Technique:
    - DL Heel Raise Up and Down together X 10 Repetitions
    - DL Heel Raise Up together, Raise Unaffected Leg and Down/Lower Affected Leg Only X 10 Repetitions
    - SL Heel Raise Affected Leg Only Up and Down X 10 Repetitions
    - DL Heel Raise Up and Down Together X 10 Repetitions
  - Duration 2 weeks
- **Phase 5:** Performed **4 days/ Week (2 days on, 2 Day off)- 1X/ Day**
  - Technique:
    - DL Heel Raise Up and Down together X 10 Repetitions
    - DL Heel Raise Up together, Raise Unaffected Leg and Down/Lower Affected Leg Only X 10 Repetitions
    - SL Heel Raise Affected Leg Only Up and Down X 10 Repetitions, 2 sets per session
    - DL Heel Raise Up and Down Together X 10 Repetitions
  - Duration 2 weeks
- **Phase 6:** Performed **3-4 days/ Week (1 days on, 1 Day off)- 1X/ Day**
  - Technique:
    - DL Heel Raise Up and Down together X 10 Repetitions
    - DL Heel Raise Up together, Raise Unaffected Leg and Down/Lower Affected Leg Only X 10 Repetitions
    - SL- Affected Leg Only Up and Down X 10 Repetitions, 3 sets
    - DL Heel Raise Up and Down Together X 10 Repetitions
  - Duration 1 month



## References

- Akizuki et al. The relative Stress on the Achilles Tendon During Ambulation in an Ankle Immobiliser: Implications for Rehabilitation After Achilles Tendon Repair. *Br J Sports Med.* 2001; 35: 329-334.
- Amin et al. Performance Outcomes After Repair of Complete Achilles Tendon Ruptures in National Basketball Association Players. *Am J Sports Med.* 2013; 41(8): 1864-1868.
- Clanton et al. Return to Play in Athletes Following Ankle Injuries. *Sports Health.* 2012; 4(6): 471-474.
- Cook, Gray. *Movement: Functional Movement Systems* (book). 2010; ON Target Publications.
- Garrison et al. The Reliability of the Vail Sport Test™ as a Measure of Physical Performance Following Anterior Cruciate Ligament Reconstruction. *Inter J Sports Phys Ther.* 2012; 7(1):20-30.
- Goulbourne, Shaun. *Sports-Related Injuries in the Young Athlete* (book from training course). 2011; Cross Country Education, LLC.
- Hamilton et al. Triple Hop Distance as a Valid Predictor of Lower Limb Strength and Power. *J Athletic Training.* 2008; 43(2): 144-151.
- Hewett et al. Understanding and Preventing ACL Injuries: Current Biomechanical and Epidemiologic Considerations- Update 2010. *N Am J Sports Phys Ther.* 2010; 5(4): 234-251.
- Huang et al. Rehabilitation Regimen After Surgical Treatment of Acute Achilles Tendon Ruptures: A Systematic Review with meta-analysis. *Am J Sports Med.* 2014; 43(4): 1008-1016.
- Kangas et al. Achilles Tendon Elongation After Rupture Repair. *Am J Sports Med.* 2007; 35(1): 59-64.
- Kangas et al. Early Functional Treatment Versus Early Immobilization in Tension of the Musculotendinous Unit After Achilles Rupture Repair: A Prospective, Randomized, Clinical Study. *J Trauma Acute Care Surg.* 2003; 6: 1171-1180.
- Maffulli et al. Early Weight-Bearing and Ankle Mobilization After Open Repair of Acute Mid-Substance Tears of the Achilles Tendon. *Am J Sports Med.* 2003. 31(5): 692-700.
- Kearney et al. The Achilles Tendon Total Rupture Score: a Study of Responsiveness, Internal Consistency and Convergent Validity on Patients with Acute Achilles Tendon Ruptures. *Health and Quality of Life Outcomes.* 2012; 10:24 1-7.
- Maffulli, N. Current Concepts Review- Rupture of the Achilles Tendon. *J Bone Joint Surg Am.* 1999; 81: 1019-1036.
- Maquirriain, J. Achilles Tendon Rupture: Avoiding Tendon Lengthening During Surgical Repair and Rehabilitation. *Yale J Bio Med.* 2011; 84: 289-300.
- McNair et al. Biomechanical Properties of the Plantar Flexor Muscle-Tendon Complex 6 Months Post-Rupture of the Achilles Tendon. *J Ortho Res.* 2013; 9999: 1-7.
- Nilsson-Helander et al. The Achilles Tendon Total Rupture Score (ATRS) Development and Validation. *Am J Sports Med.* 2007; 35(3): 421-425.

- Padua et al. The Landing Error Scoring System (LESS) is a Valid and Reliable Clinical Assessment Tool of Jump-Landing Biomechanics: The Jump-ACL Study, *Am J Sports Med.* 2009; 10:1996-1002.
- Power et al. Delayed Recovery of Velocity-Dependent Power Loss Following Eccentric Actions of the Ankle Dorsiflexors. *J Appl Physiol.* 2010; 109: 669-676.
- Reiman, Michael P., Manske, Robert C. *Functional Testing in Human Performance: 139 tests for sport, fitness, and Occupational Settings* (book). 2009. Human Kinetics.
- Rettig et al. Potential Risk of Rerupture in primary Achilles Tendon Repair in Athletes Younger Than 30 Years of Age. *Am J Sports Med.* 2005; 33(1): 119-123.
- Ross MD, Langford B, Whelan PJ. Test-retest Reliability of 4 Single-Leg Horizontal Hop Tests. *J Stength Cond Res.* 2002; 4:617-622.
- Saxena, A, Ewen, B, Maffulli, N. Rehabilitation of the Operated Achilles Tendon: Parameters for Predicting Return to Activity. *J Foot Ankle Surg.* 2011; 50: 37-40.
- Strom, AC, Casillas, MM. Achilles Tendon Rehabilitation. *Foot Ankle Clin N Am.* 2009; 14: 773-782.
- Suchak et al. Postoperative Rehabilitation Protocols for Achilles Tendon Ruptures: A Meta-analysis. *Clin Ortho Rel Res.* 2006; 445: 216-221.